

## **INTRODUCTION-**

*Granulomatous mastitis is an inflammatory condition of non-infectious origin that specifically affects breast tissue. Its etiology is unknown. Granulomatous mastitis generally occurs in young women, often after a breast-feeding period, and within a period of 5 years following childbirth.*

## **CASE REPORT**

*Our patient was a G2P1L1, previous LSCS. Had history of lump excision for. Left breast 2 years back. Presented to us at 34 weeks of gestation with 12 x10cm multilobulated mass, tense and tender. After not responding to antibiotics and incision and drainage, Histopathology showed epithelioid granuloma with multiloculated giant cells. Started on Anti tubercular therapy. Patient was taken up for ELECTIVE LSCS at 37 weeks with INCISION and DRAINAGE of left breast abscess under General anaesthesia.*



## **INTRA OP FINDINGS**

*Horizontal incision made at 12'o clock position, 50ml pus drained sent for pus c/s. Breast tissue chunks sent for MGIT, GeneXpert, bacterial and Fungal culture. Of which all the reports were found to be negative.*



## **REFERENCES**

- Barreto DS, Sedgwick EL, Nagi CS, Benveniste AP. Granulomatous mastitis: etiology, imaging, pathology, treatment, and clinical findings. Breast cancer research and treatment. 2018 Oct;171:527-34.
- Liu G, McRitchie D, Russell E, Cates EC. Spontaneous, granulomatous mastitis in a pregnant patient: A case report and review of the literature. Heliyon. 2023 Nov 1;9(11).
- Sheybani F, Sarvghad M, Naderi H, Gharib M. Treatment for and clinical characteristics of granulomatous mastitis. Obstetrics & Gynecology. 2015 Apr 1;125(4):801-7

## **DISCUSSION**

*IGM is a rare condition that is difficult to treat due to limited data and lack of treatment consensus. Early core biopsies for microbiology and histology should be performed in inflammatory breast masses with atypical appearances (large solid component, sinus formation or fistulation) or after a short period of failed antimicrobial therapy (1–2 weeks) A multidisciplinary pathway involving breast surgeons, radiologists, pathologists, infectious diseases specialists and a specialty with expertise in prescribing azathioprine (rheumatology or gastroenterology) is recommended*

## **CONCLUSION**

*Incisional drainage and antibiotic therapy, rather than steroids and surgery, are useful for treating granulomatous mastitis during pregnancy. The condition can also be effectively managed during delivery. Abstinence from breastfeeding and steroidal therapy after delivery have also proven to be effective in managing this disease. The use of steroids during pregnancy should, however, be treated with caution, taking their side effects into consideration.*